



## The "red" decline of Norway spruce or "røde rødgraner" - is it ammonium overload or top-dying?

Engvild, K.C.

*Publication date:*  
2005

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Engvild, K. C. (2005). *The "red" decline of Norway spruce or "røde rødgraner" - is it ammonium overload or top-dying?* Denmark. Forskningscenter Risoe. Risoe-R No. 1513(EN)

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Risø-R-1513(EN)

# Risø-R-Report

## The “Red” Decline of Norway Spruce or “Røde Rødgraner” – Is it Ammonium Overload or Top-Dying?

Kjeld C. Engvild

Risø National Laboratory  
Roskilde  
Denmark  
April 2005

**Author:** Kjeld C. Engvild  
**Title:** The “Red” Decline of Norway Spruce or “Røde Rødgraner” – Is it Ammonium Overload or Top-Dying?  
**Department:** Biosystems Department

**Risø-R-1513(EN)**  
**April 2005**

**Abstract (max. 2000 char.):**

**ISSN 0106-2840**  
**ISBN 87-550-3430-6**

In 1989 a new disease “røde rødgraner” or “red” decline of Norway spruce (*Picea abies*) became a serious problem in plantations in Jutland on poor, sandy soils. Some trees became red and lost their needles over a few years. The reddening started from the shoot tips.

The only important pollutant in mainly rural Jutland is ammonia from the many farms, resulting in N depositions of 10-30 kg ha<sup>-1</sup> year<sup>-1</sup>. The ammonium hypothesis of Nihlgård (1985) stipulates that certain types of forest decline is caused by an overload of ammonia from the air. Hence the possibility that “røde rødgraner” was caused by nitrogen overload was investigated. A sensitive biochemical indicator of nitrogen status of conifer trees is the content of free amino acids of the urea cycle, arginine and ornithine. In pot experiments with four times overload of N as ammonium nitrate the free arginine content increased more than 100 times, up to almost 2 % of the fresh weight. Also trees suffering from phosphorus or potassium deficiency showed large increases in arginine.

**Contract no.:**  
EU DG VI Project 93.60.DK0050  
SN232-0030

**Group's own reg. no.:**  
1310117

**Sponsorship:**

**Cover :**

Needle samples from “røde rødgraner” (taken from the 6<sup>th</sup> whorl at minimum metabolism in December) showed normal arginine and ornithine contents. There was no indication of nitrogen overload. The “red” Norway spruce may suffer from “top-dying” a common disorder of Norway spruce in Great Britain, believed to be caused by several mild winters in a row. In that case the symptoms should diminish after the very cold winter 1995-96.

**Pages:** 16  
**Tables:** 5  
**References:** 38

Risø National Laboratory  
Information Service Department  
P.O.Box 49  
DK-4000 Roskilde  
Denmark  
Telephone +45 46774004  
[bibl@risoe.dk](mailto:bibl@risoe.dk)  
Fax +45 46774013  
[www.risoe.dk](http://www.risoe.dk)

# Contents

**Introduction 4**

**Materials and Methods 5**

**Results 7**

**Discussion 10**

Acknowledgments 12

**References 13**

## Introduction

Since 1989 Norway spruce (*Picea abies*) in Danish plantations, especially on poor, sandy soils in western Jutland has been seriously damaged with a complex of symptoms which in Denmark is called "røde rødgraner" or "red" Norway spruce (Yde-Andersen 1991, Saxe & Larsen 1992, Saxe 1993). This syndrome where the needles redden and die from the shoot tips inwards have resulted in more than 10 % dead trees in certain places and much premature logging. Hypotheses on the possible causes are numerous: air pollution, acid rain, N-overload from the air, drought, nutrient leaching, potassium deficiency or other nutrient imbalances, salt damage, and mild winter damage, and sums of accumulated stresses. The disease seems not to be caused by pathogens or pests. Most work in Denmark has been concerned with three hypotheses:

- 1) Air pollution with ammonium (Nihlgård 1985)
- 2) Mild winter damage
- 3) Combined stresses

Most damage has occurred far away from densely populated areas and removed from major sources of industrial and traffic pollution. The only important pollutant in the area is ammonia liberated from pig and dairy farming with deposition rates of 10-30 kg ha<sup>-1</sup> nitrogen (Asman et al. 1998, Andersen and Hovmand 1995). One case of ammonium damage is known in chlorotic Scots pine (*Pinus sylvestris*) in the Netherlands where ammonia deposition rates are much higher than in Denmark (Van Dijk and Roelofs 1988, Pearson and Stewart 1993). Saxe (1993) analysed the contents of inorganic nutrients in "red" Norway spruce; he did not see any significant nutrient imbalance. It is known that the content of free amino acids is a sensitive measure of nutrition imbalances and a number of other environmental factors (Durzan and Stewart 1967, 1983, Stewart and Larher 1980, Gezelius and Näsholm 1993).

The purpose of this work is to establish if amino acids could be used as biochemical indicators for overdoses of nitrogen in Norway spruce (Wild and Schmitt 1995). Such work has been done for other *Picea* species (Durzan and Stewart 1967, 1983). The work will focus on the amino acids arginine and ornithine, members of the urea cycle (Aarnes et al. 1995) which are very important in amino acid metabolism and storage in the gymnosperms.

## Materials and Methods

Two year old Norway spruce (*Picea abies*) were obtained from the Tree Improvement Station, Krogerupvej 21, 3050 Humlebæk, Denmark. The original seeds were from Silkeborg (Denmark) and Westerhof (Northern Germany). A preliminary experiment was cultivated with 4-year old clones made from cuttings.

The seedlings were grown in open air for two seasons in 10 L pots in very pure quartz sand, fit for glass fabrication. Ten plants were grown per pot. There were four replications per treatment. The pots were placed over 10 L reservoir flasks and equipped with an automated air-lift watering system with drainage back into the reservoir (Haahr 1975). The plants were watered twice a day for two hours. During winter the plants were kept around 5°C in a greenhouse. The basal nutrient solution was designed for Norway spruce with the relations of N:K:P:Mg:Ca:S of 100:50:16:5:5:9 and 0.5 g nitrogen per pot added as ammonium nitrate (Ingestad 1979). Higher nitrogen (2, 4, and 8 fold) was given in two ways, either as additional ammonium nitrate (0.5, 1.5, 3.5g N), or as urea (0.5, 1.5, 3.5 g N) painted on the needles as a 10 % solution to simulate ammonium deposition from the atmosphere. All fertilisers except nitrogen were added at the start of the growing season. Nitrogen was given three times during shoot development from April till the end of June. All water used was deionized.

Samples for amino acids were taken during the winter to minimize diurnal and other variations associated with changes in growth phases (Durzan et al. 1983, Kim *et al* 1995). Current year needle samples from an observation plot for "red" Norway spruce decline at Feldborg skovdistrikt (also observed by Saxe 1993) were taken from the south and east branches of the 6th whorl. The samples were transported at 0-1°C and kept at -18 °C until analysis.

Amino acids were extracted by disintegrating 200 mg (Ultraturrax) of current year needles and 0.4 g polyvinylpyrrolidone with an internal standard of norleucine in 10 ml of modified Redgwell's mixture (Redgwell 1980) of methanol:CHCl<sub>3</sub>:water 12:5:3. The homogenate was centrifuged, and the precipitate reextracted with 10 ml of methanol:water 1:4. The combined supernatants were separated in two phases by adding chloroform and water, and the green chloroform phase discarded. The extracts were rotoevaporated to dryness and taken up in 1 ml water. The solution was passed through 10 ml sephadex C-25 cation exchange gel (H<sup>+</sup>) form, washed with 30 ml of H<sub>2</sub>O, and the amino acids eluted with 30 ml 0.2 M NH<sub>3</sub>. The ammonia eluate was lyophilized, the residue taken up in 0.3 ml water, transferred to a dram glass and dried in vacuo. The amino acids were converted to phenylisothiocyanate derivatives according to the picotag method and analysed on a Merck-Hitachi high pressure liquid chromatograph on a 30 cm C-18 column using an acetate buffered acetonitrile gradient with EDTA (Cohen and Strydom 1988).

Four similar experiments were run. Each experiment lasted two seasons where the plants were kept in the greenhouse at 2-8 °C (as low as possible) during the winter between two seasons. An experiment was started every year for 4 years. The first experiment was preliminary, involving only N overload, using quite large trees from cuttings. The plan of the other three experiments can be read from Table 1. Trees were grown without nitrogen; with Ingestad balanced nutrients; with 2, 4 and 8 times N excess either as ammonium nitrate in the nutrient solution or as urea painted on the needles; with phosphorus deficiency, potassium deficiency, sulphur deficiency, calcium

deficiency, or magnesium deficiency; or with plants that received only nitrogen as ammonium nitrate or urea, but none of the other nutrients. The experiments also involved the micronutrients, but the results are not included because they were not different from the balanced nitrogen control.

Growth analyses involved fresh weight, dry weight, height and the needle dry weight in percent of the total above ground dry weight in order to obtain a quantitative measure of the needle loss after N overload. Only fresh weight and needle percentage are reported in the present paper.

## Results

Table 1 gives the growth responses and arginine and ornithine contents of the four year old plants measured after two seasons of growth. There are very large differences

*Table 1 Growth, arginine and ornithine contents in the first large pot experiment. The small trees harvested after 2 seasons, 4 years old. Arginine increased up to more than 2 per cent of the fresh weight under large N overload. Ornithine increased to 0.3 percent.*

| Treatment                      | Fresh weight<br>Gram plant <sup>-1</sup> | Arginine<br>Gramkg <sup>-1</sup><br>needles | Ornithine<br>Gramkg <sup>-1</sup><br>needles |
|--------------------------------|--|---|--|
| No nitrogen added              | 2.8 ± 0.4                                | 0.07 ± 0.02                                 | 0.008 ± 0.004                                |
| Balanced nitrogen (Ingestad)   | 11.6 ± 0.4                               | 0.10 ± 0.04                                 | 0.009 ± 0.006                                |
| 2 x nitrogen, ammonium nitrate | 18.7 ± 1.6                               | 3.8 ± 1.1                                   | 0.10 ± 0.03                                  |
| 4 x nitrogen, ammonium nitrate | 16.8 ± 1.5                               | 15.7 ± 2.6                                  | 2.2 ± 0.5                                    |
| 8 x nitrogen, ammonium nitrate | 19.1 ± 2.5                               | 23.7 ± 3.8                                  | 3.1 ± 0.6                                    |
| 2 x nitrogen, urea on needles  | 17.3 ± 1.2                               | 1.7 ± 0.7                                   | 0.016 ± 0.007                                |
| 4 x nitrogen, urea on needles  | 15.0 ± 1.8                               | 14.5 ± 4.0                                  | 1.1 ± 0.5                                    |
| 8 x nitrogen, urea on needles  | 12.6 ± 2.5                               | 20.4 ± 2.1                                  | 2.9 ± 0.7                                    |
| Minus phosphorus               | 3.7 ± 0.3                                | 19.0 ± 1.5                                  | 0.39 ± 0.04                                  |
| Minus potassium                | 6.4 ± 0.7                                | 4.2 ± 1.3                                   | 0.15 ± 0.05                                  |
| Minus sulfur                   | 10.8 ± 0.8                               | 0.18 ± 0.04                                 | 0.013 ± 0.003                                |
| Minus calcium                  | 9.7 ± 0.2                                | 0.13 ± 0.07                                 | 0.015 ± 0.008                                |
| Minus magnesium                | 10.6 ± 0.8                               | 0.21 ± 0.07                                 | 0.006 ± 0.002                                |
| + N, minus all other nutrients | 4.4 ± 1.2                                | 17.9 ± 4.7                                  | 0.56 ± 0.25                                  |
| + N as urea, minus all others  | 2.7 ± 0.3                                | 16.3 ± 1.6                                  | 0.56 ± 0.35                                  |

between treatments, but also very large variation in the genetically inhomogenous material. Even trees in pots minus nitrogen or + nitrogen minus all other nutrients continue growing during both seasons. The differences in arginine and ornithine contents are very large, more than 200 fold. At the highest levels more than 2 % of the needle fresh weight consists of arginine.

Table 2 gives examples of total free amino acid analyses on plants grown on the control (Ingestad balanced nitrogen), on 8 times N excess, and with phosphorus deficiency. Besides the very large increases in arginine, ornithine and lysine under both



*Table 2. Amino acids in Norway spruce needles grown for two years on Ingestad nutrient solution (control), with 8 times N-overload as ammonium nitrate or with phosphorus deficiency. Large increases in arginine, ornithine and lysine.*

| Amino acid    | 8 x N overload<br>$\mu\text{moles g}^{-1} \text{ fw}$ | Control<br>$\mu\text{moles g}^{-1} \text{ fw}$ | - phosphorus<br>$\mu\text{moles g}^{-1} \text{ fw}$ |
|---------------|---|--|---|
| Arginine      | 136.0   | 0.60   | 109.0   |
| Ornithine     | 23.0  | 0.07   | 2.9   |
| Lysine        | 5.0   | 0.04   | 3.3   |
| Asparagine    | 6.7   | 0.03   | 0.38  |
| Proline       | 3.2   | 0.15   | 0.80  |
| Glutamine     | 2.6   | 0.08   | 0.55  |
| Aspartic acid | 0.73  | 0.19   | 0.64  |
| Glutamic acid | 0.70  | 0.66   | 1.09  |
| Serine        | 0.32  | 0.26   | 0.25  |
| Glycine       | 0.02  | 0.11   | -   |
| Citrulline    | 0.02  | 0.01   | 0.01  |
| Threonine     | 0.79  | 0.05   | 0.19  |
| Alanine       | 0.30  | 0.17   | 0.23  |
| Tyrosine      | 0.15  | 0.05   | 0.06  |
| Valine        | 0.17  | 0.07   | 0.08  |
| Methionine    | 0.18  | 0.04   | 0.08  |
| Cysteine      | -   | 0.06   | 0.01  |
| Isoleucine    | 0.12  | 0.04   | 0.07  |
| Leucine       | 0.13  | 0.04   | 0.08  |
| Phenylalanine | 0.20  | 0.13   | 0.13  |

N excess and P deficiency there are large increases in asparagine, glutamine and proline under N overload.

Table 3 gives the results on arginine contents in the four different experiments. The results of the N overloading results in 130-160  $\mu\text{moles g}^{-1}$  in all four experiments,

*Table 3. Arginine contents compared in the four different experiments after 2 seasons of growth.*

|                        | Control                             | 8 x N overload                      | - phosphorus                        |
|------------------------|-------------------------------------|-------------------------------------|-------------------------------------|
| Treatment              | $\mu\text{moles g}^{-1} \text{ fw}$ | $\mu\text{moles g}^{-1} \text{ fw}$ | $\mu\text{moles g}^{-1} \text{ fw}$ |
| Preliminary experiment | 0.41                                | 162                                 | -                                   |
| First experiment       | 0.60                                | 136                                 | 109                                 |
| Second experiment      | 1.4                                 | 163                                 | 19                                  |
| Third experiment       | 1.5                                 | 146                                 | 66                                  |

corresponding to 2.2-2.8 % of arginine in the needles. The phosphorus deficiency experiments have given more variable results.

The contents of arginine tend to build up over two seasons as shown in table 4. There is no response to potassium deficiency after the first season, but some response after the second season. The response to N excess and P deficiency is strong already after the first season, but increases even more after the second season.

*Table 4. Arginine contents measured after the first and second season growth in pots.*

|                       | First season             | Second season            |
|-----------------------|--------------------------|--------------------------|
| Treatment             | $\mu\text{moles g}^{-1}$ | $\mu\text{moles g}^{-1}$ |
| Control (Ingestad)    | 0.7                      | 1.5                      |
| 8 times overload urea | 52                       | 115                      |
| Minus phosphorus      | 19                       | 67                       |
| Minus potassium       | 1.2                      | 4.1                      |

Table 5 shows the ratio of needles dryweight to dryweight of the above ground biomass, expressed in percent. This is only a rough measure of needle loss, but it can be seen that N overloaded trees have fewer needles than the controls. This is because these trees had lost almost all second and third year needles.

Table 5. Reduction in the amount of needles in N-overloaded trees, expressed as needles in percent of total dry weight.

| Treatment                            | Needles, percent of dry weight |
|--------------------------------------|--------------------------------|
| Control (Ingestad nutrients)         | 46 $\pm$ 3                     |
| 8 times overload N, ammonium nitrate | 33 $\pm$ 3                     |
| 8 times overload N, urea             | 31 $\pm$ 3                     |
| Minus phosphorus                     | 37 $\pm$ 3                     |

Figure 1 shows the arginine contents in 50 trees suffering from "red" decline of Norway spruce, compared with small trees overloaded with nitrogen. The arginine and ornithine content of the damaged trees falls within the normal range. There is no indication of the arginine and ornithine increases seen after N-overload or nutrient imbalance.

## Discussion

Basically, the present work on the relation between nutrition and amino acid content confirms and extends previous results. But it does differ from the results of the many previous studies in two important ways: First, the effects of nitrogen excess or lack of potassium and phosphorus are much more dramatic in the present long-term, "chronic" pot experiments than in most other reports, except those of Durzan and Stewart (1967) and Gezelius and Näsholm (1993). The amounts of free arginine + ornithine reached more than 2 percent of the needle fresh weight under phosphorus deficiency or severe nitrogen excess. The levels obtained correspond to those found by van Dijk and Roelofs (1988) in ammonium poisoned pines in the Netherlands. Second, the study is a link between the short term work in water cultures or pots in growth chambers or greenhouses (Durzan and Stewart 1967, 1983, Gezelius and Näsholm 1993, Kim et al. 1987, van Dijk *et al* 1990), ammonia fumigation (Pérez-Soba et al. 1994), pollution investigations involving ammonia (Van Dijk and Roelofs 1988, Huhn and Schulz 1996), and the studies on amino acids circulation in xylem (Rennenberg et al. 1998, Schneider et al. 1996) and long term forest fertiliser experiments (Aronsson 1985, Edfast et al. 1990).

The amounts of free amino acids at severe nitrogen excess or phosphorus deficiency are so large that the plant cells must be in an abnormal osmotic state. (Table 1, Figure 1). Even at a modest nitrogen excess, two times N there was 0.2-0.4 % free arginine present after two growth seasons.

The results on arginine, ornithine and other amino acid contents were quite variable; the variance on sampling exceeded the variance on the chemical analyses. If one includes the large seasonal variations observed by e. g. Durzan and Stewart (1983) and Kim and Glerum (1995) it is obvious that arginine and other amino acid contents may well be sensitive biochemical indicators of nutrient imbalances, but they are not

very precise. High levels of arginine and ornithine indicate unbalanced nutrition, but not exactly what is wrong.




Figure 1. Arginine content in needles of Norway spruce suffering from the “red” decline or “røde rødgræner” syndrome and needle loss, compared with arginine content in needles from small trees overloaded with N as ammonium nitrate in pot cultures.

We did not see arginine and ornithine accumulation induced by deficiency of sulphur, calcium, magnesium (table 1) or the micronutrients (data not shown). It is possible that these nutrients have no influence on the free amino acid contents; but it is also possible that the 2 year old plants that were used for the experiments had already accumulated sufficiently large reserves of these compounds to last for several growing seasons.

An important part of the experiments was the amino acid contents of the dying Norway spruce in the Danish plantations in Jutland compared to the N-overloaded pot experiments. In the observation plot of the spruce in "red" decline in December 1995 there were 13 trees with mild damage (10-25% needle loss), 30 trees with medium damage (26-60% needle loss) and 7 trees with heavy damage (61-99% needle loss). About 10 trees were dead (markings has disappeared on many dead trees).

A major part of the results are given in figure 1. When one compares the low arginine contents of Norway spruce trees in "red" decline with the ammonium nitrate controls, there is no indication at all that the sick and dying trees suffer from major nutrient imbalance. This agrees with the work of Saxe (1993) who analysed the same trees for inorganic nutrients and saw no indication of nutrient imbalance. This contradicts our initial assumption that the mortality could be explained by the "ammonium hypothesis" of Nihlgård (1985). The results of van Dijk and Roelofs (1988) have shown that one will reach arginine levels above 100  $\mu\text{moles g}^{-1}$  in the needles of Scots pine in areas with heavy ammonium loads in the Netherlands and that these levels are found in diseased, chlorotic trees. Healthy Scots pine had levels below 30  $\mu\text{moles g}^{-1}$  arginine.

If the "red" decline in Norway spruce is not caused by ammonium, which is believed to be the major pollutant in rural Jutland, what might then be the cause? Barklund (1983) believe that many different factors contribute to damage and mortality: drought, acid rain causing aluminium toxicity, frost, and facultative parasites.

Another very simple hypothesis (Engvild 1996, 1998) is that the disease is identical to "top-dying" (Saxe and Larsen 1992). In Great Britain this physiological disorder actually limits the cultivation of Norway spruce (Diamandis 1978a, b, c, 1979 a, b, Murray 1953). This disease is characterised by a "spectacular browning or reddening ..... the needles redden from the shoot tips ..... the trees die from the top down over 2-3 years" (cited from Phillips and Burdekin, 1982). These symptoms are close to the ones described by Saxe (1993) and Saxe and Larsen (1992). "Top-dying" is believed to be caused by several mild, windy winters in a row, perhaps worsened by spring drought (Diamandis 1979b, Phillips and Burdekin 1982). In Denmark we have had 8 mild, windy winters in a row since 1987. The first "red" decline appeared in 1989.

If the "red" decline is indeed identical to "top-dying" of Norway spruce, then the symptoms should disappear after a couple of cold winters. "Red" Norway spruce decline may thus be moved from the field of pollution to the field of plant pathology (Kandler and Innes 1995 ) In the last years the symptoms of "red" spruce have been much less prevalent along with several winters with snow.

## Acknowledgments

This work was supported by EU DG VI grant 93.60.DK0050 to the National Forest and Nature Agency, Denmark. The author is grateful to J. C. Briand Petersen and S Fjederholt of Feldborg skovdistrict for machine- and manpower help in sampling the trees in "red" decline. The author is grateful to Liselotte Meltotte for the amino acid analyses.

## References

- Aarnes H, Eriksen A B and Southon T E (1995) Metabolism of nitrate and ammonium in seedlings of Norway spruce (*Picea abies*) measured by in vivo  $^{14}\text{N}$  and  $^{15}\text{N}$  NMR spectroscopy. *Physiol. Plant.* 94, 384-390.
- Andersen H V and Hovmand M F (1995) Ammonia and nitric acid dry deposition and throughfall. *Water Air Soil Pollut.* 85, 2211-2216.
- Asman W A H, Sutton M A and Schjørring J K (1998) Ammonia: emission, atmospheric transport and deposition. *New Phytol.* 139, 27-48.
- Aronsson A (1985) Indikationer på stress vid obalans i trädens växtnäringsinnehåll. *K. Skogs- Lantbr. Akad. Tidskr. Suppl.* 17, 40-51.
- Barklund P (1983) Topptorka på äldre og yngre gran. Tänkbara orsaker: torka, frost sur nederbörd, svamp. *Sveriges Skogsvårdsförbunds Tidskrift* 81 (2), 29-40.
- Cohen S A and Strydom, D J (1988) Amino acid analysis utilizing phenylisothiocyanate derivatives. *Anal. Biochem.* 174, 1-16.
- Diamandis S (1978a) "Top-dying" of Norway spruce, *Picea abies* (L.) Karst. with special reference to *Rhizosphaera kalkhoffii* Bubák. I. Development of foliar symptoms. *Eur. J. For. Path.* 8, 337-345.
- Diamandis S (1978b) "Top-dying" of Norway spruce, *Picea abies* (L.) Karst. with special reference to *Rhizosphaera Kalkhoffii* Bubák. II. Status of *R. kalkhoffii* in "top-dying" of Norway spruce. *Eur. J. For. Path.* 8, 345-356.
- Diamandis S (1978c) "Top-dying" of Norway spruce, *Picea abies* (L.) Karst. with special reference to *Rhizosphaera Kalkhoffii* Bubák. III. Moisture content of current and second year needles of Norway spruce. *Eur. J. For. Path.* 8, 357-361.
- Diamandis S (1979a) "Top-dying" of Norway spruce, *Picea abies* (L.) Karst. with special reference to *Rhizosphaera kalkhoffii* Bubák. IV. "Top-dying" of Norway spruce and climate. *Eur. J. For. Path.* 9, 78-88.
- Diamandis S (1979b) "Top-dying" of Norway spruce, *Picea abies* (L.) Karst. with special reference to *Rhizosphaera kalkhoffii* Bubák. VI. Evidence related to the primary cause of "top-dying". *Eur. J. For. Path.* 9, 183-191.
- Durzan D J and Steward, F C (1967) The nitrogen metabolism of *Picea glauca* (Moench) Voss and *Pinus banksiana* Lamb. as influenced by mineral nutrition. *Can. J. Bot.* 45, 695-710.
- Durzan D J and Steward, F C (1983) Nitrogen metabolism. In *Plant physiology a Treatise*. Vol 8: Nitrogen metabolism. Ed F C Steward. pp. 55-265. Academic Press, Orlando, Fl. ISBN 0-12-668608-4.
- Edfast A-B, Näsholm T and Ericsson A (1990) Free amino acid concentrations in needles of Norway spruce and Scots pine trees on different sites in areas with two levels of nitrogen deposition. *Can. J. For. Res.* 20, 1132-1136.
- Engvild K C (1996) Sidste år med "røde rødgraner"? *Skoven* 1996 (8), 356-357.
- Engvild K C (1998) The ammonium hypothesis: arginine content and the "red" decline or "top dying" of Norway spruce. *Ambio* 27, 158-159.

- Gezelius K and Näsholm T (1993) Free amino acids and protein in Scot pine seedlings cultivated at different nutrient availabilities. *Tree Physiol.* 13, 71-86.
- Haahr V (1975) Comparison of manual and automatic irrigation of pot experiments. *Plant Soil* 43, 497-502.
- Huhn G and Schulz H (1996) Contents of free amino acids in Scots pine needles from field sites with different levels of nitrogen deposition. *New Phytol.* 134, 95-101.
- Ingestad T (1979) Mineral nutrient requirements of *Pinus silvestris* and *Picea abies* seedlings. *Physiol. Plant.* 45, 373-380.
- Kandler O and Innes J L (1995) Air pollution and forest decline in central Europe. *Environ. Pollut.* 90, 171-180.
- Kim Y T and Glerum C (1995) Seasonal free amino acid fluctuations in red pine and white spruce needles. *Can. J. For. Res.* 25, 697-703.
- Kim Y T, Glerum C, Stoddart J and Colombo S J (1987) Effect of fertilization on free amino acid concentrations in black spruce and jack pine containerized seedlings. *Can J. For. Res.* 17, 27-30.
- Murray J S (1954) Two diseases of spruce under investigation in Great Britain. *Forestry* 27, 54-62.
- Nihlgård B (1985) The ammonium hypothesis - an additional explanation to the forest dieback in Europe. *Ambio* 14, 2-8.
- Pearson J and Stewart G R (1993) The deposition of atmospheric ammonia and its effects on plants. *New Phytol.* 125, 283-305.
- Pérez-Soba M, Stulen I and van der Eerden L J M (1994) Effect of atmospheric ammonia on the nitrogen metabolism of Scots pine (*Pinus sylvestris*) needles. *Physiol. Plant.* 90, 629-636.
- Phillips D H and Burdekin D A (1982) Top dying of Norway spruce. *In: Diseases of Forest and Ornamental Trees.* pp. 134-136. MacMillan, London. ISBN 0 333 32357 2.
- Redgwell R J (1980) Fractionation of plant extracts using ion-exchange sephadex. *Anal. Biochem.* 107, 44-50.
- Rennenberg H, Kreutzer K, Papen H, and Weber P (1998) Consequences of high loads of nitrogen for spruce (*Picea abies*) and beech (*Fagus sylvatica*) forests. *New Phytol.* 139, 71-86.
- Saxe H (1993) Triggering and predisposing factors in the "Red" decline syndrome of Norway spruce (*Picea abies*). *Trees* 8, 39-48.
- Saxe H and Larsen J B (1992) "Røde rødgraner" - økofysiologiske aspekter. *Dansk Skovbrugs Tidsskrift* 77, 187-205.
- Schneider S Gessler A, Weber P, von Sengbusch D, Hanemann U and Rennenberg H (1996) Soluble N compounds in trees exposed to high loads of N: a comparison of spruce (*Picea abies*) and beech (*Fagus sylvatica*) grown under field conditions. - *New Phytol.* 134: 103-114.
- Stewart G R, Larher F (1980) Accumulation of amino acids and related compounds in relation to environmental stress. *In The Biochemistry of Plants A comprehensive*

- treatise. Eds: Stumpf P K, Conn E E. Vol. 5 Amino Acids and Derivatives. Ed: Miflin B J. pp. 609-635.
- van Dijk H F G, de Louw M H J, Roelofs J G M and Verburgh J J (1990) Impact of artificial, ammonium-enriched rainwater on soils and young coniferous trees in a greenhouse. part II. Effects on the trees. *Environ. Pollut.* 63, 41-59.
- van Dijk H F G and Roelofs J G M (1988) Effects of excessive ammonium deposition on the nutritional status and condition of pine needles. *Physiol. Plant.* 73, 494-501.
- Wild A and Schmitt V (1995) Diagnosis of damage to Norway spruce (*Picea abies*) through biochemical criteria. *Physiol. Plant.* 93, 375-382.
- Yde-Andersen N A (1991) Røde rødgraner. *Skoven* (1) 26-28.



## **Mission**

To promote an innovative and environmentally sustainable technological development within the areas of energy, industrial technology and bioproduction through research, innovation and advisory services.

## **Vision**

Risø's research **shall extend the boundaries** for the understanding of nature's processes and interactions right down to the molecular nanoscale.

The results obtained shall **set new trends** for the development of sustainable technologies within the fields of energy, industrial technology and biotechnology.

The efforts made **shall benefit** Danish society and lead to the development of new multi-billion industries.